



AMERICAN JOURNAL OF **BIOSCIENCE AND BIOINFORMATICS (AJBB)**

VOLUME 2 ISSUE 1 (2023)



PUBLISHED BY
E-PALLI PUBLISHERS, DELAWARE, USA

Trends of Oat Production Area, Productivity and Utilization Status in Ethiopia

Lensa Urgesa Akasa^{1*}

Article Information

Received: March 20, 2023

Accepted: April 02, 2023

Published: April 10, 2023

Keywords

*Animal Feed, Ethiopia,
Household Consumption,
Oat Production, Trends*

ABSTRACT

Oat is an important multi-purpose cereal crop cultivated for grain, feed, straw used for human consumption and livestock feed. Oat is ranked as sixth in the world's cereal production following wheat, maize, rice, barley and sorghum. Therefore, this study is initiated to analyze the trends of; oats production area, productivity and utilization status in Ethiopia. Data of meher season oats area of production (ha), productivity (qt/ha) and utilization (%) were collected from the Central Statistical Agency (CSA) of Ethiopia. Trend test was carried out using the non-parametric Mann-Kendall's trend test packaged in XLstat. This study's result indicated that oat production showed a non-significant increasing trend. Additionally oats yield qt/ha showed that significant increasing trend. Oat production area showed significant decreasing despite being used as Household Consumption and animal feed showed non-significant decreasing trends.

INTRODUCTION

Oat (*Avena sativa* L.) is one of the well-adapted and important fodder crops grown in the highlands of Ethiopia, mainly under rain-fed conditions (Amanuel *et al.*, 2019) and in Ethiopia used for human consumption and livestock feed. Oat is an important multi-purpose cereal crops cultivated for grain, feed and straw over more than 9 million hectares globally (FAO, 2011). Gebremedhin *et al.* (2015) reported that oats grain is the staple diet of human beings in some parts of the central high lands of Ethiopia. Lulseged (1987) also stated that oat is a dual purpose forage crop which can also be used as human food in many parts of the world. According to the Ethiopia Ministry of agriculture and rural development, oats was registered as grass and used as forages (Getnet, 2012). Oats are abundantly grown in the central highlands of Selale and some parts of west Shewa like Meta-Robi and Galessa areas of Dendi woreda, Arsi, Bale and Gojjam. Oat is ranked as sixth in the world's cereal production following wheat, maize, rice, barley and sorghum. The initial aim of oats introduce to the smallholders was for feed production (Fekadu *et al.*, 2018).

Oat is forage crop grown at medium to high altitudes (1600-3000m) on heavy soils (vertisols) where temperate grasses and other improved forages are difficult to establish (Negash *et al.*, 2017). Oats were early maturing, palatable, succulent and energy-rich crop and mostly used as silage and is preferred by animals due to its high palatability and softness. Its grain is also a valuable feed for dairy cows, horses, young breeding animals and poultry. Many cultivars of oat have high feed value if cut at its 50 percent flowering stage which is the best time for the crop harvest for better yield and can meet the demand of the rapidly growing livestock industry of Ethiopia (Gebremedhin *et al.*, 2015).

Ethiopia is endowed with diverse agro-ecologies suitable for cereal production. Cereals crop like Oats productions in Ethiopia were under rain fed production system. However, the success of this production system relies more on climate condition. The facts about land use, utilization and number of householder at the national level have potentially imposed limits on the areas sown as a major source of increase in production. This causes inverse relationship between area and production because expansion area which is unsuitable for agriculture and uses. The interconnection of oat production area, productivity and utilization change with areas cover on trends of oat is less understood. The current trends of oat production areas, productivity and utilization have not yet investigated in Ethiopia. Therefore, the objective the study was designed to analysis trend of oat production areas, utilization in Ethiopia.

METHODOLOGY

Data of maher season for oat production area, productivity, utilization status of oat for the period of 2008 to 2020/2021 were collected from Central Statistical Agency (CSA). Trend test were carried out using the non-parametric Mann-Kendall's trend test which is less sensitive to outliers and test for a trend in a time series without specifying whether the trend is linear or non-linear (Partal and Kahya, 2006; Yenigun *et al.*, 2008; Hadgu *et al.*, 2013). The Mann-Kendall's test statistic is given as:

$$S = \sum_{i=1}^{N-1} * \sum_{j=i+1}^N \text{sgn}(x_j - x_i) \quad \text{Equation (1)}$$

Where S is the Mann-Kendal's test statistics; x_i and x_j are the sequential data values of the time series in the years i and j ($j > i$) and N is the length of the time series. A positive S value indicates an increasing trend and a negative value indicates a decreasing trend in the data series.

¹ Oromia Agricultural Research Institute, Mechara Agricultural Research Center, P. O. Box 19, Machara, Ethiopia

* Corresponding author's e-mail: lensa Urgesa15@gmail.com

The sign function is given as

$$\text{sgn}(x_j - x_i) = \begin{cases} +1 & \text{if } (x_j - x_i) > 0 \\ 0 & \text{if } (x_j - x_i) = 0 \\ -1 & \text{if } (x_j - x_i) < 0 \end{cases} \quad \text{Equation (2)}$$

The variance of S , for the situation where there may be ties (i.e., equal values) in the x values

$$\text{var}(S) = \frac{1}{18} [N(N-1)(2n+5) - \sum_{i=1}^m t_i(t_i-1)(2t_i+5)] \quad \text{Equation (3)}$$

Where, m is the number of tied groups in the data set and t_i is the number of data points in the i th tied group. For n larger than 10, Z_{MK} approximates the standard normal distribution (Partal and Kahya, 2006; Yenigun *et al.*, 2008) and computed as follows

$$Z_{MK} = \begin{cases} \frac{S-1}{\sqrt{\text{var}(S)}} & \text{if } S > 0 \\ 0 & \text{if } S = 0 \\ \frac{S+1}{\sqrt{\text{var}(S)}} & \text{if } S < 0 \end{cases} \quad \text{Equation (4)}$$

The presence of a significant trend is evaluated using the Z_{MK} value. In a two-sided test for trend, the null hypothesis H_0 should be accepted if $Z_{MK} < Z_{1-\alpha/2}$ at a given level of significance. $Z_{1-\alpha/2}$ is the critical value of Z_{MK} from the standard normal table.

RESULTS AND DISCUSSION

Sen's slope value in Tables 1 to 3 indicated that the country-level production of Oats (Qt/yr) showed a non-significant ($p=0.879$) increasing trend by factors of 3313Qt

for 2008-2020 production years. In the country Yield of Oat Qt/ha showed significant ($P=0.000$) increasing trend by factor of 0.823 (Qt/ha)/year. However, Oat production area showed significant ($P=0.041$) decreasing trend by factor of -778.695 ha/year. In agreement with this result, (Wasihun and Desu, 2020) indicated that in Ethiopia area of oat production is decrease despite yield Qt/ha is increase. Household Consumption showed a non-significant ($P=0.087$) decreasing trend by factor of -0.186 percent/year. Oat used as animal feed showed a non-significant ($P=0.542$) decreasing trend by factor of -0.027 percent/year.

As CSA show that oat production is distributed only in Oromia, Amhara and SNNP. From One decade ago Oat is abundantly grown in the central highlands of Selale and some parts of west Shewa like Meta-Robi and Galessa areas of Dendi worda, Arsi, Bale and Gojjam. However, recently due to the expansion of crop cultivation, the area of oat production is decreased and oat used as household consumption and animal feed is decreased while cereal and pulse crop production area is increased and household consumption is more shifted to cereal and pulse crops.

But production of oat (Qt/Yrs) and yield (Qt/ha) of oat is increased, this is due to various research result which is related to oat variety development, agronomy and adaption with modern inputs. Generally in Ethiopia due to shortage of land, the attentions of small householders are on cash crops and human feed crops rather than animal feed.

Table 1: Oat production (Qt), Production area (ha) and Yield (Qt/ha) in Ethiopia from 2004-2020

Years	Production in(Qt)	Production area (Ha)	Yield (Qt / Ha)
2004	566,754.00	45,131.00	12.56
2005	401,634.00	44,401.00	9.05
2007	365,857.00	30,556.00	11.97
2008	427,729.00	30605.00	13.98
2009	330,191.00	24,017.00	13.75
2010	475,650.57	30,858.76	15.41
2011	494,749.24	30,568.39	16.18
2012	436,337.83	26514.10	16.46
2013	616,502.59	35,617.76	17.31
2014	508,059.26	27,899.64	18.21
2015	402,689.43	22,105.72	18.22
2016	491,796.43	24,040.94	20.46
2017	526,318.93	25,896.22	20.32
2018	301,439.40	14,843.08	20.31
2019	457,543.61	21,281.80	21.5
2020/21	381,530.36	21,882.01	19.7

Source CSA, 2004-2020/21

Table 2: Oat Utilization in Ethiopia from 2008-2020

Years	Household Consumption %	Oat used for Animal feed %
2008	65.39	1.37
2009	67.62	0.28
2011	59.39	0.85
2012	62.32	0.95
2013	60.67	1.05
2014	67.02	1.17
2015	64.19	0.69
2016	60.99	0.86
2017	58.86	0.58
2019	61.11	0.67
2020/21	52.73	0.89

Source, CSA, 2008-2020/21

Table 3: Mann-Kendall Trend statistics for Production, Area of Production, Yield, and Household Consumption, Animal Feed for 2008-2020/21.

Oats	Zmk	MKStatistics	p-value	Alpha	Sen's slope:	Trend
Production area (ha)	-0.491	-27.000	0.041	0.05	-778.695	Decrease
Production (Qt)	0.055	3.000	0.879	0.05	3313	Increase
Yield (Qt/ha)	0.818	45.000	0.000	0.05	0.823	Increase
Household consumption (%)	-0.418	-23.000	0.087	0.05	-0.816	Decrease
Animal feed (%)	-0.164	-9.000	0.542	0.05	-0.027	Decrease




Figure 1: Production of Oats in Ethiopia from 2003-2020




Figure 2: Production area of Oat in Ethiopia from 2004-2020




Figure 3: Productivity (Yield Qt/ha) of Oat in Ethiopia from 2004-2020




Figure 4: Oat used as Household Consumption in Ethiopia from 2008-2020




Figure 5: Oats used as Animal feed (%) in Ethiopia from 2008-2020

CONCLUSION

Oat is an important multi-purpose cereal crop cultivated for grain, feed and straw used for human consumption and livestock feed. Sen's slope value in Tables 1 to 3 indicated that the country-level production of Oat (Qt/yr) showed a non-significant ($p=0.879$) increasing trend by factors of 3313Qt for 2008-2020 production year. In the country yield of Oat Qt/ha showed significant

($P=0.000$) increasing trend by factor of 0.823 (Qt/ha)/year. However, Oat production area showed significant ($P=0.041$) decreasing trend by factor of -778.695 ha/year. Household Consumption showed a non-significant ($P=0.087$) decreasing trend by factor of -0.186 percent/year. Oat used as animal feed showed a non-significant ($P=0.542$) decreasing trend by factor of -0.027 percent/year for 2008-2020 production year.

REFERENCE

- [illegible]

- Republic of Ethiopia Central Statistical Agency, Agricultural Sample Survey (2009 E.C.), volume I, report on area and production of major crops (private peasant holdings, meher season) statistical bulletin 584. Addis Ababa, Ethiopia.
- Central Statistical Agency (2016). The Federal Democratic Republic of Ethiopia Central Statistical Agency, Agricultural Sample Survey (2009 E.C.). Volume vii, report on crop and livestock product utilization (private peasant holdings, meher season) statistical bulletin 586. Addis Ababa, Ethiopia
- Central Statistical Agency (2017). The Federal Democratic Republic of Ethiopia Central Statistical Agency, Agricultural Sample Survey (2010 E.C.), volume I, report on area and production of major crops (private peasant holdings, meher season) statistical bulletin 586. Addis Ababa, Ethiopia.
- Central Statistical agency (2017). The Federal Democratic Republic of Ethiopia Central Statistical Agency, Agricultural Sample Survey (2010 E.C.). Volume vii, report on crop and livestock product utilization (private peasant holdings, Meher season) statistical bulletin 588. Addis Ababa, Ethiopia
- Central Statistical Agency (2018). The Federal Democratic Republic of Ethiopia Central Statistical Agency, Agricultural Sample Survey (2011 E.C.), volume I, report on area and production of major crops (private peasant holdings, meher season) statistical bulletin 589. Addis Ababa, Ethiopia.
- Central Statistical Agency (2019). The Federal Democratic Republic of Ethiopia Central Statistical Agency, Agricultural Sample Survey (2012 E.C.), volume I, report on area and production of major crops (private peasant holdings, meher season) statistical bulletin 587. Addis Ababa, Ethiopia.
- Central Statistical Agency (2019). The Federal Democratic Republic of Ethiopia Central Statistical Agency, Agricultural Sample Survey (2010 E.C.). Volume vii, report on crop and livestock product utilization (private peasant holdings, meher season) statistical bulletin 588. Addis Ababa, Ethiopia
- Central Statistical Agency (2020). The Federal Democratic Republic of Ethiopia Central Statistical Agency, Agricultural Sample Survey (2013 E.C.). Volume vii, report on crop and livestock product utilization (private peasant holdings, meher season) statistical bulletin 592. Addis Ababa, Ethiopia
- Central Statistical Agency (2020). The Federal Democratic Republic of Ethiopia Central Statistical Agency, Agricultural Sample Survey (2013 E.C.), volume I, report on area and production of major crops (private peasant holdings, meher season) statistical bulletin 590. Addis Ababa, Ethiopia.
- Gebremedhn B, Araya HA, Gebremedhn BH. (2015). Evaluation of different oat varieties for fodder yield and yield related traits in Debre Berhan Area, Central Highlands of Ethiopia. *Livestock Research for Rural Development* 27(9).
- Getnet Assefa, Mesfin Dejene, Jean Hanson, Getachew Anemut, Solomon Mengistu and Alemayehu Mengistu. (2012). Forage Seed Research and Development in Ethiopia. Ethiopian Institute of Agricultural Research. Addis Ababa, Ethiopia.
- Hadgu G, Tesfaye K, Mamo G, Kassa B. (2013). Trend and variability of rainfall in Tigray, Northern Ethiopia: Analysis of meteorological data and farmers' perception. *Academically Journal Environmental Science* 1(8), 159-171.
- Mosissa, F, Kefala, B., & Abeshu, Y. (2018). Potential of oats (*avena sativa*) for food grain production with its special feature of soil acidity tolerance and nutritional quality in central highlands of ethiopia. *Advances in Crop Science and Technology*, 6(4), 1-5.
- Negash D, Animut G, Urgie M, Mengistu S. (2017). Chemical Composition and Nutritive Value of Oats (*Avena sativa*) Grown in Mixture with Vetch (*Viciavillosa*) with or Without Phosphorus Fertilization in East Shoa Zone, Ethiopia. *J Sci Food Agric*, 87(5), 89-108.
- Partal T, Kahya E. (2006). Trend analysis in Turkish precipitation data. *Hydrological.Processes*, 20(9), 2011-2026.
- Wasihun Gizaw and Desu Assegid. (2020). Trend of cereal crops production area and productivity, in Ethiopia.
- Wada, A., Shawle, K., & Gemiyo, D. (2019). Biomass yield and nutritional quality of different oat varieties (*Avena sativa*) grown under irrigation condition in Sodo Zuriya Distric, Wolaita Zone, Ethiopia. *Agricultural Research*, 20(4), 1-10.
- Yenigün K, Gümüş V, Bulut H. (2008). Trends in stream flow of the Euphrates basin, Turkey. *Proclamation Institute of Civil Engineering Water Management*, 161(4), 189-198.