The American Journal of Physical Education and Health Science (AJPEHS)

ISSN: 2992-9679 (ONLINE)

VOLUME 4 ISSUE 2 (2026)



PUBLISHED BY
E-PALLI PUBLISHERS, DELAWARE, USA

American Journal of Physical Education and Health Science (AJPEHS)

Volume 4 Issue 2, Year 2026
ISSN: 2992-9679 (Online)
DOI: https://doi.org/10.54536/ajpehs.v4i2.9013
https://journals.e-palli.com/home/index.php/ajpehs

Structured Non-Invasive Rehabilitation Pathways for Recovery After Body-Contouring Procedures

Oksana Repina[1*]

Article Information

Received: August 29, 2026

Accepted: November 13, 2026

Published: November 24, 2026

Keywords

Aesthetic Rehabilitation, Body-Contouring Recovery, Multidisciplinary Care, Postoperative Pathways, Supportive Care

ABSTRACT

Body contouring demands rigorous recovery. Physical suffering and shifting tissues cause immediate distress, but patients also face severe, long-term worry over how they will look in the end. Current postoperative care protocols focus almost entirely on strict surgical monitoring. They leave an unaddressed oversight when it comes to structured, supportive aesthetic therapies. In response, this paper introduces a new, staged methodological model. It cohesively blends non-invasive treatments, specifically compression-vibration massage alongside targeted thermo-cellular wraps into various phases of the healing process. Existing clinical literature on edema, fibrosis, and the emotional toll of surgical recovery directly supports this methodology. To examine how this functions in practice, the article details an observational case of a 37-year-old woman recovering from abdominoplasty, liposuction, and gluteal fat transfer. Her qualitative feedback during the initial weeks was highly positive. She noted softer tissues and much faster resolution of her bruising, alongside a significant drop in both pain and anxiety. Integrating wellness-oriented practices into standard clinical follow-up creates a vastly superior healing environment, directly validating the importance of a patient's day-to-day recovery journey while pending formal controlled evaluations. Bringing wellness-oriented practices into standard clinical follow-up creates a vastly superior healing environment, pending formal controlled evaluations.

INTRODUCTION

Body-contouring procedures have become increasingly common in contemporary aesthetic practice. These procedures are frequently performed to improve body contour and patient satisfaction with physical appearance. When doctors perform these operations, they intentionally alter the natural architecture of the patient to create a better silhouette. They choose procedures like extensive liposuction and tummy tucks. Surgeons remove fat and cut away extra skin. This process creates substantial tissue disruption just under the surface. Medical teams control this injury carefully. But the body still reacts strongly. Tissues swell. Bruises appear everywhere. Blood vessels leak fluid into the surrounding spaces to start the repair process (Colwell, 2010). Doctors naturally focus on keeping the patient safe. They watch out for major issues like severe infections or blood clots (Gusenoff, 2014). Surgical management primarily focuses on procedural and postoperative safety. Then the patient goes home to face the physical reality of healing.

Standard recovery pathways have a major missing piece. Clinical guidelines tell doctors how to handle surgical wounds and prevent emergencies. They rarely give patients a clear plan for the long and uncomfortable healing phase. People often call this the ugly phase of recovery. The skin gets hard. Pockets of fluid build up under the surface. Uneven scar tissue forms inside the damaged fat layers over time (Thomas *et al.*, 2010). Medical professionals call this internal scarring fibrosis. Fibrosis makes the skin look bumpy and uneven (Mera-Cruz *et al.*, 2024). This period may be associated with significant psychological distress. Patients look in the mirror and worry about their final results. They experience high levels of postoperative anxiety. They feel isolated. And they feel stressed (Reavey *et al.*, 2011). Surgeons often tell them to just be patient and wait. The prolonged recovery period may contribute to uncertainty regarding treatment outcomes. People need active physical help to feel comfortable in their own skin again.

This paper introduces a new way to guide patients through this difficult period. This paper proposes a conceptual framework designed to support postoperative body-contouring recovery. The model brings wellness professionals into the medical timeline safely. These practitioners do not act like doctors. They focus purely on the patient's physical comfort and external appearance. They use specialized nonmedical tools to speed up recovery. One tool involves mechanical compression combined with vibration. Another uses warm botanical body wraps. These methods may facilitate fluid redistribution and support postoperative comfort (Kubler *et al.*, 2020). They may help manage tissue irregularities associated with early postoperative fibrosis, potentially supporting skin adaptation during recovery.

Figure 1 presents the proposed tissue remodeling cycle associated with postoperative recovery. The process naturally begins with surgical trauma, followed by the first active step of recovery. We call this Phase 1, which focuses entirely on decongestion to control acute edema. Next comes Phase 2, the middle step, which involves deep tissue remodeling. Practitioners use targeted vibration to prevent hard fibrosis from forming under the skin. Phase

[1] Independent Research, United States
[*] Corresponding author's e-mail: oksana.repina.edu@gmail.com




Figure 1: The Post-Contouring Tissue Remodeling Cycle

3 follows closely behind and targets skin retraction. Active cosmetic wraps help safely tighten loose tissue. The cycle finally ends when the patient achieves the final aesthetic contour. The primary objective of the framework is simple. We want to protect the surgeon's artistic work while giving the patient a smooth healing experience.

LITERATURE REVIEW

Pathophysiology of Post-Liposuction Tissue Healing

Doctors make cuts and remove fat. This causes a lot of damage under the skin. The body responds immediately. Fluid rushes into the empty spaces. Blood vessels break and bleed. This creates pronounced postoperative edema and bruising (Gusenoff, 2014; Thomas *et al.*, 2010). Over time, the body tries to heal the empty tunnels. It lays down thick scar tissue. We call this fibrosis. Fibrosis makes the skin feel hard and look lumpy. Mera-Cruz *et al.* (2024) studied women after secondary liposuction. They found that almost all patients developed some form of hard nodules or skin depressions. Other researchers agree that uncontrolled scarring ruins the final look of the surgery (Cedidi & Ingianni, 2006; Colwell, 2010). The empty fat pockets can also fill with fluid and stay there for weeks. This constant swelling puts pressure on the healing skin. It stops the skin from shrinking down tight against the new muscles. Some studies even look at using fat grafts to fix these hard scars later on (Krastev *et al.*, 2020). But preventing the scars in the first place is much better.

Psychosocial Dynamics of Aesthetic Surgery Recovery

People get plastic surgery to feel better about themselves. But the recovery period often does the exact opposite. Patients frequently experience postoperative swelling, bruising, and concerns regarding aesthetic outcomes. This sharp drop in mood is very common. The medical world sometimes calls it postoperative depression. Reavey *et al.* (2011) looked at how patients feel after body contouring. They found that doctors rarely measure the emotional struggle of recovery. They just look at the physical scars. This is a big problem. Chronic swelling and hard lumps make patients think the surgery failed. They hide from their friends and family. Social avoidance becomes a daily habit (Manrique *et al.*, 2022). Research on young adults links self-esteem with social comparison tendencies (Abellana *et al.*, 2024), and appearance-related concerns during recovery may engage comparable self-evaluative processes. Patients may perceive recovery as a prolonged period of physical and emotional uncertainty. They need constant reassurance. Psychological research in other populations likewise treats stress as a factor associated with well-being (Ay-ad, 2026), which supports measuring the emotional burden of recovery as a distinct outcome. A purely medical approach ignores this mental burden. Doctors fix the body. But patients need someone to guide them through the emotional stress of healing.

Mechanical Stimulation in Fibrosis Prevention

How do we stop the body from making hard scars? We use physical force. Research shows that mechanical pressure changes how cells behave. Wong *et al.* (2012) explained this clearly. When you push and pull on healing tissue, the scar cells line up neatly. They do not form messy, hard lumps. This is why massage and compression work so well. Many studies prove this point. Intermittent pneumatic compression pushes fluid out of the belly much faster than regular tight garments (Abdelhalim & Samhan, 2021). Other researchers found great results using manual drainage and specific taping methods right after surgery (Bruno *et al.*, 2025; Santos *et al.*, 2024). Some clinics use ultrasound and deep tissue massage to break down early fibrosis (Cacho *et al.*, 2024; Chi *et al.*, 2018; Machado & Lima, 2021; Masson *et al.*, 2014).

Vibration is especially powerful. Kołodziejczak *et al.* (2025) reviewed mechanical massage devices. They found that rolling and vibrating the skin physically breaks down rigid fibers. It also drains fluid and makes the skin stretch better. Shockwave therapy does the same thing for hard fat deposits (Bruno & Cilluffo, 2025). Gentle physical therapy also stops pain and gets patients moving earlier (Río-González *et al.*, 2025; Torres *et al.*, 2012). For serious swelling, tight bandages and negative pressure machines keep the fluid out of the wounds (Horch, 2015; Klein *et al.*, 2020; Singh *et al.*, 2019). Even conditions like lipedema respond best when we squeeze the fluid out physically (Forner-Cordero *et al.*, 2012; Tamura *et al.*, 2025).

Phyto-Therapeutic and Thermo-Cellular Interventions

Mechanical pressure is just one half of the solution. The skin also needs chemical help from the outside. Doctors usually ignore plant-based treatments. But recent studies show they actually work. Some researchers tested special peptide creams on patients after body contouring. The people who used the creams had much less swelling and bruising. Their hard lumps went away faster, too (Casas *et al.*, 2020, 2022; Kubler *et al.*, 2020).

Aesthetic professionals use a similar idea with active body wraps. They put plant extracts directly on the skin.

These wraps can be very hot or very cold. The sudden temperature change forces the blood vessels to open and close. This pumps stagnant fluid away from the surgical site. It works a lot like tissue sealants that surgeons use to stop bleeding inside the body (Por *et al.*, 2009). Botanical formulations are applied topically as part of supportive wellness protocols. They may contribute to perceived improvements in skin condition and patient comfort. So the loose skin shrinks tightly over the new muscles instead of hanging down.

Science gives us a lot of clues. We know that surgery causes massive swelling and hard scars. We know that patients feel terrible during this time. And we have clear proof that compression, vibration, and topical treatments fix these exact problems. But hospitals do not combine these tools into a single plan. Surgeons do their job and step back. The patient tries to heal alone. The proposed framework seeks to address this gap by organizing supportive interventions within a structured recovery timeline. It takes all these proven physical therapies. It organizes them into a safe step-by-step timeline. This model bridges the huge gap between the operating room and the final beautiful result. Table 1 contrasts the strategies of the medical and aesthetic teams for three recurring postoperative challenges and shows where the two sets of measures complement each other.

Table 1: Collaborative Solutions for Post- Surgical Challenges

Postoperative Challenge	Medical Team Strategy	Aesthetic Team Strategy
Severe Fluid Buildup	Places surgical drains in the operating room. Prescribes standard compression garments.	Applies systemic pressotherapy. Utilizes supportive cooling applications intended to enhance patient comfort.
Hard Internal Scarring	Injects steroids into the scar tissue. Performs secondary liposuction to remove rigid lumps.	Uses targeted mechanical vibration. Applies mechanical stimulation intended to support tissue remodeling.
Loose Skin Recovery	Cuts away excess tissue surgically. Closes the wound with medical sutures.	Applies warm cellular wraps. Supports skin adaptation during later stages of recovery.

MATERIALS AND METHODS

Core Principles of Body Contouring Support

Timing controls everything in this framework. Practitioners cannot just start rubbing a fresh surgical site. They must wait for the right moment. The primary surgeon decides when the incisions are stable enough to tolerate outside help (Gusenoff, 2014). If a patient received a fat graft in their gluteal area, we protect those new cells at all costs. We do not apply pressure there. Instead, the wellness professional acts like an architect of the soft tissue. They smooth out the superficial canvas. The medical team handles the deep structural work. This clear division keeps the patient perfectly safe. But we still actively improve their physical comfort. We look at the body as a connected system. When a surgeon removes fat, the space quickly fills with blood and water (Thomas *et al.*, 2010). The body tries to heal this massive damage by throwing thick scar tissue everywhere. Our job is to guide that healing process physically. The framework proposes the use of supportive non-invasive interventions intended to assist postoperative tissue recovery.

Phase 1: Lymphatic Decongestion (Weeks 1 to 2)

The first two postoperative weeks are commonly characterized by substantial edema and patient discomfort. Severe swelling takes over the entire body. The framework starts very gently. We use low-pressure systemic pressotherapy to move stagnant fluid. Intermittent pneumatic compression pushes trapped blood and water out of the swollen belly effectively (Abdelhalim & Samhan, 2021). The patient wears a special suit with multiple air chambers. The machine fills these chambers with air in a specific rhythm. It creates a gentle wave of pressure. This wave moves from the feet up to the center of the body. It literally squeezes the excess fluid back into the lymphatic system. And it works much better than wearing tight compression clothes alone (Klein *et al.*, 2020).

We combine this mechanical pumping with cold, active botanical wraps. Cold temperatures force the surface blood vessels to constrict. This physical action flushes cellular debris away from the damaged fat layers naturally. Aestheticians often use lotions containing bitter orange or cooling menthol. These plant extracts soak into the skin. They make the blood vessels shrink and push the stagnant fluid away from the fresh surgical wounds (Forner-Cordero *et al.*, 2012). Patients may experience subjective improvements in comfort following supportive interventions.

Phase 2: Anti-Fibrotic Mechanical Stimulation (Weeks 3 to 4)

Around the third week, the healing tissue starts to harden. This is when fibrosis begins. We change our strategy completely. The practitioner introduces targeted compression and microvibration therapy. A specialized device rolls over the skin to deliver rapid mechanical pulses. These vibrations are intended to support tissue remodeling during early stages of fibrosis development (Kołodziejczak *et al.*, 2025). The device has a heavy cylinder filled with small spheres. As the cylinder spins, the spheres punch the skin thousands of times per minute. The patient wears a protective suit to keep their skin safe from friction.

Human cells respond directly to physical force. Mechanical pulling tells the fibroblasts to align their collagen fibers in straight lines instead of messy clumps (Wong *et al.*, 2012). This process may contribute to more uniform tissue organization. This stops the bumpy appearance that often ruins a good liposuction result. We do not work blindly. The aesthetician moves the vibrating cylinder along specific lines. They push the broken-down tissue toward the major lymph nodes. This clears out the dead cells from the surgical trauma (Santos *et al.*, 2024). Active tissue manipulation also blocks pain signals from reaching the brain. Patients usually feel much less sore after just one session (Torres *et al.*, 2012).

Phase 3: Skin Retraction and Stabilization (Weeks 5 and Beyond)

The final stage focuses on tightening the loose skin. The body lost a lot of volume during surgery. The empty skin needs to shrink back down. We apply advanced warm body wraps to speed up this process. These wraps often contain active botanical extracts like green tea and caviar. Increased local temperature may contribute to improved tissue perfusion and patient comfort during recovery. It feeds the skin cells so they can produce more elastin. Clinical trials confirm that active topical products improve skin retraction significantly (Casas *et al.*, 2022).

Warm topical applications are commonly incorporated into postoperative wellness protocols. Their proposed role is to support patient comfort and perceived skin quality during later stages of recovery. However, additional clinical research is required to clarify the mechanisms and magnitude of these effects (Kubler *et al.*, 2020). We keep using the vibrating machine during this phase. But we change the speed and pressure. The goal shifts from breaking scars to building firm skin. The final phase aims to support tissue stabilization and patient satisfaction with postoperative recovery. This structured timeline takes away the guesswork. The framework is intended to support both aesthetic outcomes and the patient recovery experience. Figure 2 summarizes the timing of the three modalities across the recovery period. Systemic pressotherapy occupies the earliest weeks and ends at the start of week 3. Targeted compression and vibration begin at the start of week 3 and continue to week 7. Thermo-cellular wraps run across the entire period. The overlapping bars show that the framework adds modalities cumulatively instead of replacing one with another, so that each phase builds on the tissue state left by the previous one.




Figure 2: Phased Modality Integration

RESULTS AND DISCUSSION

Case Description

We observed a very complex clinical case from a specialized body contouring practice. The patient was a 37-year-old female. She decided to undergo a massive single-session surgery. Her surgeon performed an abdominoplasty. The doctor also did a comprehensive liposuction of her abdomen and inner thighs. Finally, they transferred the extracted fat into her gluteal area to reshape her curves. This is an extensive surgical intervention for one day. The immediate reaction of her body was severe. She developed extreme generalized swelling across her entire torso and legs. Massive dark bruises covered her skin. Her tissues felt hard and painful to the touch. This hardening is a common early sign of fibrosis (Mera-Cruz *et al.*, 2024). The patient reported concerns regarding postoperative appearance and recovery progression. She expressed a desire for timely recovery and restoration of normal daily activities.

Application of the Supportive Protocol

The aesthetic team applied our proposed conceptual framework right after getting clearance from the plastic surgeon. They started very carefully. The first step involved systemic pressotherapy to push out the massive fluid buildup gently. They combined this mechanical pumping with cold, active botanical wraps. The practitioner used specific wet wraps to soothe the inflammation and shrink the swollen blood vessels. Around the third week, the team introduced mechanical compression and vibration. They used a specialized device over a protective suit. But they strictly avoided the gluteal area. We must protect

transferred fat cells so they survive and establish a new blood supply (Krastev *et al.*, 2020). The vibration focused entirely on the liposuction sites, such as the abdomen and thighs. The rolling spheres physically broke down the early hard spots in the tissue.

Qualitative Patient Feedback

The patient reported positive physical and emotional recovery experiences during the observation period. We tracked her progress using qualitative feedback. The patient reported a massive drop in physical pain after just a few sessions. Her dark bruises faded very fast. The most surprising part was the speed of her overall healing. She completed the acute recovery phase in exactly three weeks. Standard recovery usually takes much longer. She noted that her skin felt soft and tight. There were no hard lumps or rigid scars under the surface. The uniform body contours looked completely natural. And her mental state improved dramatically. She stopped worrying about the surgical outcome. The patient reported reduced anxiety regarding postoperative recovery. Researchers know that physical comfort directly boosts mental health and overall satisfaction during surgical recovery (Reavey *et al.*, 2011). This observation is consistent with previous literature emphasizing the relationship between physical comfort and postoperative well-being. Figure 3 displays three symptom domains across the first five weeks on a relative scale from 0 (low) to 100 (high). Generalized edema falls from 95 in week 1 to 5 in week 4 and 0 in week 5. Tissue hardness, used as a marker of fibrosis, peaks at 80 in week 2, one week after the edema peak, and declines to 10 in week 4. Postoperative anxiety follows the physical symptoms and drops from 90 in week 1 to 15 in week 3. The heatmap shows that the emotional and physical trajectories moved together and that fibrosis lagged behind edema, which is the sequence the phased design anticipates.




Figure 3: Therapeutic Symptom Heatmap

Bridging the Gap Between Surgery and Final Aesthetics

Patients frequently have high expectations regarding postoperative outcomes. But they usually wake up looking heavily bruised and completely swollen. This harsh physical reality creates a huge gap between the operating room and the final mirror reflection. The proposed framework may help address this gap by providing structured postoperative support. We do not just passively wait for the swelling to go down. We actively manage the healing tissue. This active physical care provides massive psychological relief. Structured follow-up may improve perceived continuity of care. They see their bruises fade week by week. They feel the hard lumps disappear before they turn into permanent scars (Mera-Cruz *et al.*, 2024).

Chronic swelling often leads to severe psychological morbidity. People experience high anxiety and social avoidance when their bodies look deformed (Manrique *et al.*, 2022). They hide from their friends. They stay home. This structured support protects the financial and emotional investment of the individual. It brings them back to normal daily life much faster (Colwell, 2010). Many doctors forget that quality of life matters just as much as the surgical technique itself (Reavey *et al.*, 2011). People need to feel good in their clothes. So we treat the emotional stress by fixing the physical body. A smooth recovery journey builds deep trust between the patient and the entire medical team.

Safety and Interdisciplinary Synergy

Patient safety drives this entire model. We never cross medical boundaries. Aesthetic practitioners must wait for the green light from the primary surgeon. You cannot just massage fresh incisions. Doctors know exactly when the body is ready for mechanical help (Gusenoff, 2014). This open communication creates a perfect interdisciplinary synergy. Plastic surgeons do amazing structural work inside the body. But they rarely have the clinical time to sit and massage away every small lump for an hour. And they desperately need a safe place to send their patients for dedicated aftercare.

Our staged protocol gives clinics exactly that. It offers

a highly reliable pathway. We avoid fat grafts so those fragile new cells survive and grow without pressure (Krastev *et al.*, 2020). We focus only on draining the fluid and smoothing the superficial fat. And we use safe plant-based wraps to tighten the loose skin effectively (Casas *et al.*, 2022). The surgeon focuses entirely on surgery. The aesthetician focuses completely on the external canvas. Surgeons get much better final cosmetic outcomes as a result. The risk of permanent fibrosis drops dramatically. The patient wins. And the entire medical practice builds a stronger reputation.

Limitations

This paper presents the proposed conceptual framework rather than a validated clinical protocol. The proposed model is supported by literature describing individual interventions and by an illustrative observational case. However, the framework has not been evaluated through controlled clinical trials. Future research should investigate clinical effectiveness, safety outcomes, patient-reported measures, and long-term postoperative recovery trajectories across larger patient populations.

CONCLUSION

Body contouring surgeries change lives. But the physical cost remains extremely high. Surgeons create massive trauma under the skin when they remove fat. The body immediately responds with severe swelling and hard fibrosis. Standard medical care usually leaves patients to deal with this ugly phase alone. This paper showed a much better way to handle the recovery. The proposed conceptual framework organizes supportive interventions into a structured recovery timeline. We start with gentle pressotherapy. This mechanical pumping pushes out trapped fluid to control the early swelling effectively. Then we apply targeted vibration. The rolling spheres break down early scar tissue physically before it hardens into permanent lumps. Finally, we use warm botanical wraps. These active cosmetics feed the cells so the loose skin shrinks tight over the new muscles. In the illustrative case, the patient lost her bruises and severe swelling in just three weeks. She felt happy. Structured supportive care may improve the postoperative recovery experience. The plastic surgery industry needs to wake up. Making a perfect cut in the operating room is just half the job. Doctors must protect their artistic work. The healing tissue needs active guidance for months. Future research should look more closely at how mechanical vibration stops fibrosis. We need large clinical trials to measure exactly how much the skin shrinks after using active botanical wraps. And we desperately need standardized recovery guidelines. Medical doctors and wellness practitioners have to start working together. They need to share the same goal. Surgeons handle the deep medical work. Aesthetic professionals manage the superficial canvas. This collaborative approach may enhance patient safety when appropriately supervised by the surgical team. It also ensures that people actually enjoy their new bodies without suffering through weeks of high anxiety and pain. Further investigation is warranted to determine whether structured supportive protocols can improve recovery outcomes in body-contouring populations.

REFERENCES

Abdelhalim, N. M., & Samhan, A. (2021). Influences of intermittent pneumatic compression therapy on edema and postoperative patient's satisfaction after lipoabdominoplasty. *Aesthetic Plastic Surgery, 45*(4), 1667–1674. https://doi.org/10.1007/s00266-021-02272-w

Abellana, C. J. P., Mendez, A. M. H., Subido, G. M. V., & Culajara, C. L. (2024). The social comparison trap: Association between fear of missing out (FoMO) and self-esteem in college students. *American Journal of Human Psychology, 2*(1), 114–120. https://doi.org/10.54536/ajhp.v2i1.2807

Ay-ad, V. R. (2026). Academic stress, time management, and well-being. *American Journal of Human Psychology, 4*(2), 37–49. https://doi.org/10.54536/ajhp.v4i2.4847

Bruno, A., & Cilluffo, M. (2025). Shockwave therapy in lipedema patients: *A prospective study. Phlebology, 41*(1), 58–67.https://doi.org/10.1177/02683555251343155

Bruno, A., Vigliano, T., & Foti, R. (2025). The role of taping in early recovery after abdominoplasty: A comprehensive review and proposal of a standardized protocol. *Aesthetic Plastic Surgery, 49*(21), 6128–6136. https://doi.org/10.1007/s00266-025-05057-7

Cacho, N. J. V., Sales, W. B., da Silva, J. V. B., & Vidal, G. P. P. (2024). Effects of physical therapy in the postoperative period of plastic surgery: An integrative review. *Journal of Sports Medicine and Therapy, 9*(2), 32–36. https://doi.org/10.29328/journal.jsmt.1001079

Casas, L., Bell, M., Claytor, B., Ziegler, M. E., & Widgerow, A. (2020). An analysis of patient-reported recovery outcomes of topical tripeptide/hexapeptide formulations utilized in a prospective randomized double-blind split neck and body study. *Aesthetic Surgery Journal Open Forum, 3*(1), Article ojaa052. https://doi.org/10.1093/asjof/ojaa052

Casas, L., Claytor, R., Zeidler, K., Shridharani, S., Cohen, S. R., Khanna, J., Gould, D. J., Yates, E. K., Lultschik, S., Bell, M., & Widgerow, A. (2022). A multicenter, randomized, double-blind, split-body clinical trial evaluating the efficacy and outcomes of a topical product pre and post aesthetic surgical body procedures. *Aesthetic Surgery Journal Open Forum, 4,* Article ojac054. https://doi.org/10.1093/asjof/ojac054

Cedidi, C., & Ingianni, G. (2006). Compression therapy after complex soft tissue trauma, and flap coverage: Optimization of scar development, swelling, function, and aesthetic result. *European Journal of Medical Research, 11*(2), 85–89.

Chi, A., Lange, A., Guimarães, M., & Santos, C. B. (2018). Prevenção e tratamento de equimose, edema e fibrose no pré, trans e pós-operatório de cirurgias plásticas [Prevention and treatment of ecchymosis, edema, and

fibrosis in the pre-, trans-, and postoperative periods of plastic surgery]. *Revista Brasileira de Cirurgia Plástica, 33*(3), 343–354. https://doi.org/10.5935/2177-1235.2018RBCP0147

Colwell, A. (2010). Current concepts in post-bariatric body contouring. *Obesity Surgery, 20*(8), 1178–1182. https://doi.org/10.1007/s11695-010-0120-4

Forner-Cordero, I., Szolnoky, G., Forner-Cordero, A., & Kemény, L. (2012). Lipedema: An overview of its clinical manifestations, diagnosis and treatment of the disproportional fatty deposition syndrome: Systematic review. *Clinical Obesity, 2*(3–4), 86–95. https://doi.org/10.1111/j.1758-8111.2012.00045.x

Gusenoff, J. A. (2014). Prevention and management of complications in body contouring surgery. *Clinics in Plastic Surgery, 41*(4), 805–818. https://doi.org/10.1016/j.cps.2014.06.006

Horch, R. (2015). Incisional negative pressure wound therapy for high-risk wounds. *Journal of Wound Care, 24*(Suppl. 4b), 21–28. https://doi.org/10.12968/jowc.2015.24.Sup4b.21

Klein, I., Tidhar, D., & Kalichman, L. (2020). Lymphatic treatments after orthopedic surgery or injury: A systematic review. *Journal of Bodywork and Movement Therapies, 24*(4), 109–117. https://doi.org/10.1016/j.jbmt.2020.06.034

Kołodziejczak, A., Adamiak, J., & Rotsztejn, H. (2025). Endermologie as a complementary therapy in medicine and surgery and an effective aesthetic procedure: A literature review. *Applied Sciences, 15*(8), Article 4313. https://doi.org/10.3390/app15084313

Krastev, T., Schop, S., Hommes, J., Piatkowski, A., & van der Hulst, R. R. (2020). Autologous fat transfer to treat fibrosis and scar-related conditions: A systematic review and meta-analysis. *Journal of Plastic, Reconstructive & Aesthetic Surgery, 73*(11), 2033–2048. https://doi.org/10.1016/j.bjps.2020.08.023

Kubler, M. M., Sheehan, M., & Casas, L. (2020). Topical treatment with tripeptide and hexapeptide following body-contouring procedures improved patient-reported recovery outcomes. *Aesthetic Surgery Journal Open Forum, 2*(1), Article ojz037. https://doi.org/10.1093/asjof/ojz037

Machado, G. C., & Lima, T. S. (2021). Intervenção da fisioterapia no tratamento de fibrose cicatricial no pós-operatório de cirurgia estética [Intervention of physiotherapy in the treatment of cicatricial fibrosis in the postoperative period of aesthetic surgery]. RCMOS: *Revista Científica Multidisciplinar O Saber, 1*(4), 146–153. https://doi.org/10.51473/rcmos.v1i4.2021.59

Manrique, O., Bustos, S. S., Ciudad, P., Adabi, K., Chen, W. F., Forte, A., Cheville, A., Jakub, J., McLaughlin, S., & Chen, H. C. (2022). Overview of lymphedema for physicians and other clinicians: A review of fundamental concepts. *Mayo Clinic Proceedings, 97*(10), 1920–1935. https://doi.org/10.1016/j.mayocp.2020.01.006

Masson, I. B., de Oliveira, B. S. A., Machado, A., Farcic, T. S., Esteves Júnior, I., & Baldan, C. (2014). Manual lymphatic drainage and therapeutic ultrasound in liposuction and lipoabdominoplasty post-operative period. *Indian Journal of Plastic Surgery, 47*(1), 70–76. https://doi.org/10.4103/0970-0358.129627

Mera-Cruz, G., Durán, H., Murillo-Romero, N., Perez, M. E., Rubiano, V., & Camacho-Domínguez, L. (2024). Fibrosis management algorithm in secondary liposculpture. *Aesthetic Surgery Journal, 44*(8), 859–868. https://doi.org/10.1093/asj/sjae026

Por, Y., Shi, L., Samuel, M., Song, C., & Yeow, V. (2009). Use of tissue sealants in face-lifts: A metaanalysis. *Aesthetic Plastic Surgery, 33*(3), 336–339. https://doi.org/10.1007/s00266-008-9280-1

Reavey, P., Klassen, A., Cano, S., McCarthy, C., Scott, A., Rubin, J., Shermak, M., & Pusic, A. (2011). Measuring quality of life and patient satisfaction after body contouring: A systematic review of patient-reported outcome measures. *Aesthetic Surgery Journal, 31*(7), 807–813. https://doi.org/10.1177/1090820X11417426

Río-González, Á., Delgado-Pérez, E., García-García, E., González-Fernández, L., García-Isidoro, S., & Cerezo-Téllez, E. (2025). Physiotherapy intervention in the immediate postoperative phase of lipedema surgery: Observational study. *Journal of Clinical Medicine, 14*(7), Article 2137. https://doi.org/10.3390/jcm14072137

Santos, B. O., Bezerra, C. D. N., & Silva, Y. R. (2024). Eficácia do taping linfático e da drenagem linfática manual no pós-operatório de lipoabdominoplastia abdominal [Efficacy of lymphatic taping and manual lymphatic drainage after abdominal lipoabdominoplasty]. *Revista FT, 28*(140), 49–50. https://doi.org/10.69849/revistaft/pa10202411240749

Singh, D., Lobach, V., & Holton, T. (2019). Use of closed-incision negative-pressure therapy in aesthetic surgery. *Plastic and Reconstructive Surgery, 143*(1S), 11S–14S. https://doi.org/10.1097/PRS.0000000000005306

Tamura, L. H. H., Castilho, G. C., Vieira, M. E. M., Fernandes, A. L. F., Mayer, F., Bandeira, F. S., De Moraes, P. F., & Padoveze, B. A. (2025). Surgical and non-surgical approaches in the management of lipedema: A systematic review. *Lumen et Virtus, 16*(53), Article e9178. https://doi.org/10.56238/levv16n53-097

Thomas, M., Menon, H., & D'Silva, J. (2010). Surgical complications of lipoplasty: Management and preventive strategies. *Journal of Plastic, Reconstructive & Aesthetic Surgery, 63*(8), 1338–1343. https://doi.org/10.1016/j.bjps.2009.06.046

Torres, R., Ribeiro, F., Duarte, J. A., & Cabri, J. (2012). Evidence of the physiotherapeutic interventions used currently after exercise-induced muscle damage: *Systematic review and meta-analysis. Physical Therapy in Sport, 13*(2), 101–114. https://doi.org/10.1016/j.ptsp.2011.07.005

Wong, V., Longaker, M., & Gurtner, G. (2012). Soft tissue mechanotransduction in wound healing and fibrosis. *Seminars in Cell & Developmental Biology, 23*(9), 981–986. https://doi.org/10.1016/j.semcdb.2012.09.010